

# An investigation of infant deaths following initial hepatitis B vaccination based on the Vaccine Adverse Event Reporting System (VAERS), 1992-2002

Valentina A. Soldatenkova<sup>1</sup>, MS; and F. Edward Yazbak<sup>2</sup>, MD, FAAP

<sup>1</sup>3282 Walnut St. #C  
Los Alamos, NM 87544  
Phone: +1 505 661 8078  
Fax: +1 505 665 4817  
Email: valyusha@mail.com

<sup>2</sup>TL Autism Research  
70 Viewcrest Drive  
Falmouth, MA 02540  
Phone: +1 508 540 5060 Fax: +1 508 457 9814  
Email: TLAutStudy@aol.com

## Abstract

**Purpose:** To examine Sudden Infant Death Syndrome (SIDS) and hepatitis B neonatal vaccination. **Results:** 170 reports related to neonatal hepatitis B vaccination were filed with the Vaccine Adverse Event Reporting System (VAERS) during 1992-2002. Of the 38 (22.4%) death reports, 29 were unexplained; 24 were classified as SIDS in the United States National Vital Statistics, and 4 were attributed to unknown causes. **Conclusion:** A systematic review of neonatal SIDS and other unexpected infant deaths following the initial dose of hepatitis B vaccination should be undertaken at the international level.

© Copyright 2007 Pearlblossom Private School, Inc.—Publishing Division. All rights reserved.

**Keywords:** SIDS, sudden death, vaccines

## 1. Introduction

Sudden *unexpected* death in infancy and early childhood was defined prior to 1969 as “the death of a child who was thought to be in good health or whose terminal illness appeared to be so mild that the possibility of a fatal outcome was not anticipated.” [1] Such definition was usually applied to children under 2 years of age [1,2]. Before 1963 and the First International Conference on Causes of Sudden Death in Infants [3], few professionals other than forensic pathologists were aware of this phenomenon.

It was only at the Second International Conference on Causes of Sudden Death in Infants in 1969 that “Sudden Infant Death Syndrome”, or SIDS, was first defined as “the sudden death of an infant which was unexpected by history and in which a thorough postmortem examination failed to demonstrate an adequate cause of death.” [4]

Until then sudden *unexpected* deaths with *known* causes were mixed with truly sudden and *unexplained* deaths [5].

Since 1989 SIDS has been defined in the United States as “the sudden death of an infant under 1 year of age which remains unexplained after a thorough case investigation, including performance of a complete autopsy, examination of the death scene, and review of the clinical history.” [6]

There was no code for SIDS in International Classification of Diseases (ICD) before 1973 [7], although related ICD-7 and ICDA-8 codes and causes of sudden *unexpected* deaths were published elsewhere [8,9]. SIDS has been housed in ICD chapter “Symptoms, Signs, and Ill-Defined Conditions” since 1973 and coded 795.0 under ICDA-8, 798.0 under ICD-9, and R95 under ICD-10. ICDA-8 and ICD-9 were used in the U.S. in 1968-78 [7] and in 1979-1998 [10], respectively. ICD-10 was implemented in the U. S. since 1999 [10].

A SIDS investigative and autopsy protocol was published and recommended for medical examiners and coroners in 1976, but was not mandated at the state and local levels. It never required review of the vaccination history [11].

In California, the SIDS protocol required an autopsy since 1974, and an inquiry about administered vaccines since 1990 [12,13].

The Centers for Disease Control and Prevention (CDC) and the National Institute of Child Health and Human Development (NICHD) released a generic protocol of the Sudden *Unexplained* Infant Death Investigation Report Form (SUIDIRF) in 1996. Only California, Minnesota, Missouri, and New Mexico had written SUID investigation protocols before 1996 [14]. Although the form required taking a vaccination history, its use was never enforced [15,16].

In 2006 the CDC reported that “some SUIDs are not investigated and, when they are, cause-of-death data are not collected and reported consistently.” [17] The CDC and other organizations revised the 1996 form and released a new Sudden *Unexplained* Infant Death Investigation (SUIDI) Reporting Form that requires taking the vaccination history on March 1, 2006 [18].

## 2. SIDS and vaccine injury compensation

In order to protect the vaccine manufacturers and health care providers, the U.S. Congress passed the National Childhood Vaccine Injury Act (NCVIA) in 1986.

Amendments to the act established the National Vaccine Injury Compensation Program (NVICP) in 1988 [19]. The Food and Drug Administration (FDA) and the CDC launched the Vaccine Adverse Event Reporting System (VAERS) in July 1990 in response to NCVIA [20].

Ridgway [21] reviewed 786 claim disputes from the start of the NVICP in 1988 through June 1996. Of the 786 claims, 107 (13.6%) resulted from DTP-related adverse events that led to early death and 73 (68.5%) of those were awarded compensation. In 50 of those 73 compensated claims, autopsy findings were interpreted as SIDS; 9 cases had “non-specific findings”, 4 had “other specific diagnosis”, 3 had pneumonia, 1 had encephalitis, and 6 were not autopsied.

Evidently the U.S. Court of Claims was convinced that the deaths were related to vaccine injury and wrongly classified as SIDS.

Only 8 infant deaths were attributed to vaccine adverse events in 1979–1996 [10], suggesting that most SIDS cases compensated through NVICP in 1988–1996 retained their SIDS code in the corresponding U.S. annual mortality files.

There were 12 *postneonatal* deaths attributed to vaccine adverse events in 1979–2002. Two deaths occurred annually in 1983, 1985, and 1999 and one in each of 1984, 1986, 1992, 1995, 2000, and 2001. [10] The underlying cause of one infant death (UCOD) in 1986 was “adverse reaction to pertussis vaccine including combinations with a pertussis component”; other conditions mentioned on the death certificate were “SIDS” and “Serum reaction other” [22].

Two often-quoted studies on SIDS and vaccines published prior to 1990 focused on DTP vaccine and SIDS beyond age 2 weeks [23] and *postneonatal* SIDS [24]. Pediatric pathologists do not consider infant deaths attributed to SIDS in the first 2 or 3 weeks of life to be “classical” SIDS suitable for SIDS research purposes [23,25]. Despite this, approximately 300 to 400 neonatal deaths were attributed to SIDS in the U. S. annually from 1979–1991 [10].

This investigation concerns neonatal SIDS and hepatitis B vaccination in reports to VAERS from the time of endorsement of the universal vaccination of newborns against hepatitis B in late 1991 [26] through the end of 2002. The Hepatitis B vaccine is the only vaccine licensed for and administered to neonates in the U.S.

*Neonatal* SIDS had declined in the U. S. from 11.9/100,000 live births in 1979 [10] to 7.0/100,000 in 1991 (Table 6).

In June 1992, the American Academy of Pediatrics (AAP) recommended placing all healthy infants in non-prone position during sleep to reduce the risk of SIDS [27]. After the implementation of the “Back to Sleep” campaign, the greatest decline in SIDS occurred among 1 to 6 month-old infants [28]. The *neonatal* SIDS rate per 100,000 live births declined from 7.3 in 1992 to 5.5 in 1996, and decline was statistically significant, but changes in rates of *neonatal* SIDS were not statistically significant in 1997–2004. Rates of *neonatal* SIDS during 1997–2004 ranged from 5.2 in 1999 to 4.6 in 2001–2003. (Table 6). The failure of neonatal SIDS to decline, concomitant with the increased use of the birth dose of hepatitis B vaccine, and the report by Silvers *et al.* [29] that some infant deaths reported to VAERS as SIDS from July 1990 to June 1997 would not have met the current SIDS case definition [6], provided the motivation for this investigation.

### 3. Materials and methods

We used VAERS data available for online search [30]. Few neonatal U.S. VAERS reports were filed prior to January 1, 1992. From January 1, 1992 to December 31, 2002, there were 170 distinct neonatal reports filed with VAERS. Of these, 38 (22.4%) were death reports. In order to determine the UCOD, these 38 reports were matched by date of death and age at death with mortality data from the U.S. National Vital Statistics (NVS).

The documentation enclosed with the main mortality files for years 1992–1994 [31–33] and the perinatal mortality files for years 1995–2002 [34–41] describes the file characteristics and the available data. We obtained available history and autopsy data for some infants from the review by Niu *et al.* [42] and from VAERS reports.

### 4. Results

Nine out of 38 deaths were due to known causes. Of 29 deaths reported as unexplained, 24 were attributed to SIDS, 4 to unknown/unspecified causes and one report remained unmatched (see Table 1, footnote “a”). Seventeen were boys out of the 29 infants whose death was “unexplained”—a male-to-female ratio of 1.4 which is similar to the male/female ratio in the highly vaccinated U. S. general population. There were approximately 1,214,556 males and 1,156,701 females born during 7 months in 1992 [10]. Since approximately 90% of SIDS deaths occur before age 7 months, and deaths of 2,836 male and 1,834 female infants under age 7 months were classified as SIDS in 1992 [31], incidence of SIDS in the first 7 months is 234 (95% C.I., 225 to 242) per 100,000 live born males and 159 (95% C.I., 151 to 166) per 100,000 live born females. The male-to-female SIDS incidence rate ratio of 1.47 (95% C.I., 1.39 to 1.56) is statistically significant (since the confidence interval does not include 1.0). Since there are no data on male-to-female SIDS incidence rate ratio in the cohort of the infant population that was not vaccinated, it is uncertain whether SIDS naturally occurs 1.5 times more among males compared to females.

Four of the 29 infants expired within 24 hours of vaccination and four others expired during the following 24 hours. Two infants who developed symptoms consistent with a vaccine reaction within hours of vaccination, were hospitalized, placed on life support, and died on day 3 and 4 post vaccination.

We extracted and reviewed all available information on county of residence of all recorded cases between 1992 and 2002. Because maternal smoking during pregnancy is a recognized risk factor of SIDS, we reviewed these data. Since perinatal mortality files and linked birth/infant death data sets were not available for years 1992–1994, information on maternal smoking was only available for years 1995–2002. In California, Indiana, South Dakota, and New York (excluding New York City) maternal smoking during pregnancy is not mentioned on birth certificates [43].

The data from the 29 VAERS reports are summarized in Tables 1 and 2. States in Tables 1 and 2 are denoted according to official U.S. postal abbreviations. Table 1 contains reports that included hepatitis B vaccine lot number. The column “Number of reports” in Table 1 contains the number of reports submitted to VAERS from each hepatitis B vaccine lot. Table 2 contains reports in which hepatitis B vaccine lot number was not reported.

#### 4.1 Lot Numbers

Cases 10, 11 and 12 received vaccines from lot 0165D. Case 11 was vaccinated on the day he was born, received no other vaccines, and died at age 55 days. Cases 2 and 3 resided in the Philadelphia (PA) City area, received a dose of hepatitis B vac-

cine from lot 1116A2 and lot 116A2, respectively. We believe these doses were from the same lot and that a recording error occurred.

Cases 20 and 21 received vaccines from lot 2930A2. This lot generated 3 reports of unexplained infant deaths, two of which were classified as SIDS and in the third, the death was attributed to “other ill-defined and unspecified causes of mortality” in NVS with ICD-10 code R99. The third infant received other vaccines concomitantly. Other hepatitis B vaccine lots in Table 1 that generated 2 or 3 infant death reports involved the same uncertainty.

#### 4.2 SIDS in New Hampshire

Silvers *et al.* [29] reported that from July 1990 through June 1997, New Hampshire had the highest VAERS reporting rate of infant deaths in any state: 2.73 reports per 10,000/year; Oregon was second with 1.16 per 10,000/year.

Of the 29 unexplained infant deaths reported after the neonatal dose of hepatitis B vaccine, 15 (51.7%) were from New Hampshire, a state with 14,041 to 17,569 live births in 1990–2002 (Table 4). This is a concern as it suggests significant underreporting from other states. Cases 8, 10, 11, 12, 13, 15, 23, 24, and 25 resided in New Hampshire, Hillsborough County.

An inquiry by one of us (FEY) to the NH Bureau of Maternal and Child Health revealed that any infant death that is referred as a possible SIDS from the Office of the Chief Medical Examiner to the NH SIDS Program is also referred to the NH Immunization Program for their follow up as a possible vaccine adverse event. Once the final diagnosis was made, a copy of the death certificate was sent to the NH Immunization Program, which then contacted VAERS from 2 months to more than a year after the infant’s death.

Tables 3 and 4 list SIDS and “infant deaths due to unknown causes” in New Hampshire during the periods 1979–1990 and 1991–2002, or twelve years before and after universal neonatal hepatitis B vaccination. Table 4 shows the number of unexplained infant deaths reported from New Hampshire to VAERS by December 31, 2002.

There were 23 unexplained neonatal deaths in New Hampshire during 1979–1990—before routine hepatitis B vaccination was recommended and the prone sleeping position discouraged. There were 25 unexplained neonatal deaths during 1991–2002, of which 12 were reported to VAERS as possible adverse outcomes of hepatitis B vaccination.

#### 4.3 Available History for Cases Listed in Tables 1 and 2

Infants born to mothers who smoked during pregnancy included cases 9, 12, 13, 15, 20, 21, 23, 24, and 25.

There were nine reports of infants co-sleeping with one or both parents during the last sleep: cases 7, 12, 13, 15, 19 through 23 [42,44]. Accidental smothering or overlying was not mentioned in any case.

Case 22 had a skull fracture (with no other information).

Three infants were found with blood around the nose: cases 3, 5, and 7. Case 7 also had some blood in the mouth [42].

Unless stated otherwise, the UCOD in the corresponding annual mortality file was SIDS.

Infant case histories presented below are ones that did not meet the definition of SIDS and/or unexplained death or had autopsy findings suggesting other diagnoses.

Case 3: This 15-day-old infant was brought to the ER in full arrest less than 48 hours following vaccination. Petechiae on the thymus, pericardium and pleura in addition to a fatty liver were seen at autopsy.

Case 5: This 20-day-old infant expired on the day following vaccination. Autopsy revealed: “Thymic cyst, right lobe; lungs: recent focus alveolar hemorrhage, focus squamous aspiration; liver: moderate microvascular changes, extramedullary hematopoiesis with focal neutrophilia; kidneys: urate deposit in collecting ducts, focal nephrosclerosis; adrenal glands: involution with focal calcification; pancreas: nesidiodyplasia; thymus: moderate eosinophil deposition.” [42]

Case 7: This infant was vaccinated at age 14 days and died 10 days later. Autopsy revealed: “cyst, edema brain, hemorrhage, lung disease”.

Case 8: This infant who was vaccinated at age 3 days, developed gastroesophageal reflux (GER) and was on ampicillin plus antibiotic eye drops. She died at age 22 days. Death was reported to VAERS as “SIDS”, but ultimately attributed to “other unknown and unspecified cause” in NVS, ICD-9 code 799.9.

Cases 12, 14, and 15 were vaccinated on the first day of life.

Case 12: This premature infant (35 weeks gestation/ 2240 g) who developed jaundice expired at age 19 days [42].

Case 13: This infant (34 weeks gestation / 1829 g) had feeding difficulties and prematurity-related apnea. She was discharged from the hospital on an apnea monitor. She was vaccinated at the age of 13 days, was hospitalized 5 days later and expired 4 days after that. She was off the apnea monitor during her last sleep, co-slept with her mother on a couch and was found supine on the floor [42].

Because preterm infants with a birth weight below 2000 g have a decreased immune response to hepatitis B vaccination, it has been recommended that vaccination be delayed until the age of 1 month. [45]. It is not known why this infant was vaccinated earlier than recommended.

Case 14: This baby with “sudden agitation, screams” and frequent vomiting episodes following vaccination expired at the age of 39 days.

Case 15: This infant developed a cold 2–3 days before death and died at age 16 days. Autopsy revealed “hepatic infarct with hyperemic border”. [42]

Case 19: A few hours after he was vaccinated at the age of 18 days, this infant became sweaty and inconsolable and the injection site appeared swollen. He was put to sleep in parents’ bed at 2:00am, and was found dead at 4:00am, 16 hours after vaccination. “SIDS” was mentioned on the report.

**Note:** There were 11 reports of ER visits following vaccination with hepatitis B vaccine lot 2612A2. One report was about a 5-year old boy from Louisiana, who received a dose from the same hepatitis B vaccine lot (and no other vaccine) on 7/20/1998, had a grand mal seizure 2 days later and 37 more seizures within 2 weeks. The child developed epilepsy and was placed on Tegretol.

Case 27: This infant had fever, vomiting, and diarrhea 2.5 hours after vaccination and improved. He was found apneic and

pulseless in the crib approximately 20 hours after vaccination. He was hospitalized and ventilated for 2 days until absence of brain activity was confirmed. The autopsy revealed “Massive myocardial infarction and diffuse ischemic encephalopathy” [42] yet the case was reported to VAERS as “SIDS”.

**Case 28:** This infant was vaccinated at age 7 days and died 4 days later. She had no congenital anomalies. The autopsy revealed both cerebral and pulmonary edema.

By December 31, 2002, there were also 30 VAERS reports listing cerebral edema after administration of hepatitis B vaccine alone or with other vaccines. 6 of the 19 individuals who died had received the recombinant hepatitis B vaccine alone.

Deaths of 3 infants were classified as SIDS in NVS: cases 7 and 28 above, and a case with VAERS ID 120390, an infant who died approximately 16 hours after the second dose of hepatitis B vaccine.

According to the SIDS histopathology atlas, cerebral edema is in the “gray zone” of SIDS, although it is a significant and potentially fatal finding by itself. [46] Was cerebral edema in all 3 cases a terminal event of a *coincidental* SIDS death, or was the death directly related to the hepatitis B vaccination that was administered 1, 4 or 10 days earlier?

**Case 29:** This jaundiced 2-day-old infant became irritable and developed grand mal seizures a few hours after vaccination. He later went into acute respiratory failure and cardiac arrest. At the hospital, he was diagnosed with acute anoxic encephalopathy, status epilepticus, acute respiratory failure and syndrome of inappropriate anti-diuretic hormone (ADH) secretion.

Of note is the fact that in cases 27 and 29, the deaths were attributed to unknown cause (ICD-9 code 799.9 and ICD-10 code R99 respectively) in NVS and not to anoxic encephalopathy.

## 5. Discussion

None of the recognized risk factors of SIDS is actually a cause of SIDS and there are no specific symptoms associated with SIDS. [7] Minor respiratory system inflammatory infiltrates, intrathoracic petechial hemorrhages, congestion and edema of the lungs are often reported autopsy findings, but are not pathognomic of the syndrome [7,47].

Beyond infancy, any death with one or more pathological findings observed in SIDS is not attributed to SIDS. The assumption that intrathoracic petechiae result from breathing against an obstructed upper airway remains theoretical and unproven because it has never been established how and where the alleged obstruction occurred [47]. Kalokerinos and Dettman have proposed that subclinical scurvy can be an alternative explanation for the intrathoracic petechiae seen in SIDS. [48] Overlooked factors requiring research include the perturbation of clotting cascade, toxic or immunological damage to the capillary basement membrane and other molecular events that accompany toxic or septic shock. [47] No effort has been made to determine the predictive value of a particular pathological finding to help confirm or exclude SIDS except for the thymic and intrathoracic petechiae [47].

Several cases had eosinophilia at various sites. Case 5 had eosinophilic deposition in the thymus. Case 12 had “chronic,

eosinophilic triaditis” of the liver and “mild peripheral eosinophilia at cerebellum/dentate nucleus and spleen”. [42]

Eosinophils are an important component of inflammation in bronchial asthma [49]. A study of 48 infants who died of SIDS and 30 who died of other, non-pulmonary, causes showed three times more eosinophils accompanied by increased T lymphocytes and B lymphocytes in the lungs of infants who died of SIDS, but the vaccination status of the study subjects was not mentioned [50].

Osborn stated in 1943 that acute hemorrhagic pulmonary edema is fatal if it persists for more than a few minutes and should be regarded as the cause of death. [51] In case 7, the site of hemorrhage was not specified in the report.

Reporting the sudden death of three infants with scurvy, Follis observed that the “liver was yellowish” and “showed atrophy of the central cells and a good deal of fatty infiltration”. [52] Fatty liver was often the only abnormal finding in Australian Aboriginal children who died suddenly and unexpectedly soon after vaccination. Such deaths were greatly reduced when Kalokerinos administered parenteral vitamin C concomitantly with the vaccine, and after vaccination [53].

Griffin *et al.* [24] reported a case of a death that occurred within 3 days of a DTP vaccination. The infant had gastrointestinal bleeding and a fatty liver and the death *was not attributed* to SIDS. Reye’s syndrome is another condition that results in fatty liver.

Oronasal bleeding [1] and frothy, blood tinged mucus secretions [11] accompany pulmonary pathology seen in both vaccinated and unvaccinated infants who have died suddenly and unexpectedly [1], but these findings are less common than thymic and intrathoracic petechiae. Kalokerinos did not observe these findings in the Australian Aboriginal children he has evaluated. [53,55].

From January 1992 through December 2002, there were approximately 60 reports of bleeding and approximately 700 reports listing liver problems filed with VAERS following Hepatitis B vaccination.

Cheraskin [54] quoting Kalokerinos [53], mentioned that smokers are likely to have lower blood ascorbate levels compared to non-smokers and suggested that additional research was needed to assess infant serum ascorbate level and its relationship to SIDS.

In this sample, six infants born to smoker mothers (cases 15, 12, 20-24) were vaccinated within the first 48 hours of life. Because the blood ascorbate levels of newborns are not routinely checked and vaccination is known to result in increased utilization of vitamin C [55], we hypothesize that those infants may have been born with relatively low level of ascorbate and an increased risk of adverse reactions to vaccination.

Silvers *et al.* [29] collected information on birth weight for children under 5 years of age whose deaths were reported to VAERS from July 1990 to June 1997, and determined that the mean birth weight of the study population (3090 g) was less than the mean birth weight of the U.S. general population (3400 g). 16.8% of the study cases were considered “low birth weight infants”(LBW). The percentages of LBW infants in the U. S. general population were 7.1% in 1991 [56] and 7.5% in 1997. [57] In our sample 2 out of 29 neonates were premature and had birth weight under 2500 g. There have been several reports of

apnea and bradycardia in some small premature babies shortly after vaccination with DTP and Hib [58-62].

There was no mention of hepatitis B immunoglobulin administration to any of the 29 neonates. This suggests that they were born to mothers who were HBsAg negative and therefore at no risk for vertical transmission of the disease. It certainly would have been safe to delay the vaccination in all these cases.

Case 15 had a minor illness before expiring. Kalokerinos reported that some of the infants he studied and who went into shock after vaccination or developed a serious illness such as pneumonia, had a minor illness when vaccinated [53,55].

The CDC has posted the following information regarding SIDS and vaccines: “The age at which infants begin their primary course of vaccinations (2 to 4 months old) is also the peak age for the incidence of sudden infant death syndrome (SIDS).” [63]

We disagree with this statement since most infants are presently vaccinated shortly after birth.

The CDC also states that: “The incidence of SIDS is declining in the United States due to public education campaigns regarding infant sleeping position, reduced exposure of infants to cigarette smoke and fewer potentially hazardous sleeping environments” and that SIDS rates decline despite the increase in vaccinations, therefore routine vaccinations do not contribute to SIDS [63].

Table 5 contains counts of U.S. resident births and infant deaths for years 1989-2004. Data from Table 5 are used to calculate mortality rates given in Table 6. The fact that both SIDS and all-cause infant mortality rates declined in 1992-1998 (Table 6) suggests that the decline in SIDS in 1992-1998 was genuine. However, all-cause *postneonatal* mortality rate reached a plateau and remained at 2.3 per 1000 live births in 1999-2002, declined to 2.2 in 2003 and returned to 2.3 in 2004. While “SIDS” continued to decline, deaths attributed to accidental mechanical suffocation and to unknown/unspecified cause increased in 1999-2002 [64,65]. The increasingly similar age at death distributions among SIDS, cause unknown/unspecified, and accidental suffocation provide *additional* evidence that there was a change in the reporting of SIDS [67].

All-cause *neonatal* mortality rate per 1000 had a statistically significant decline from 4.7 in 1999 to 4.5 in 2001 (Table 6). The 2002 increase in neonatal mortality rate was attributed to increase in the number of less than 750-gram and 500-gram live births and the high mortality among such extremely immature infants. [66] Decrease of neonatal mortality in 2004 was significant compared with neonatal mortality in 2002, but not significant in comparison with neonatal mortality in 2000, 2001, and 2003 (Table 6).

From 1989 to 1991, SIDS (among infants aged <28 days) comprised 6.4% (1051/16400; 95% C.I., 6.0% to 6.8%) of the cases. From 1999 to 2001, SIDS in this same age group comprised 8.1% (599/7405; 95% C.I., 7.5% to 8.7%). (Table 5) The incidence rate ratio of the 1999-2001 rate to the 1989-1991 rate is 1.26 (95% C.I., 1.14 to 1.40) indicating a statistically significant increase in proportion of neonatal SIDS since implementation of universal vaccination of newborns against hepatitis B and gradual increase in coverage by the birth dose.

Several epidemiological studies [23,24,67-70] concluded that vaccinations are *not a risk factor* of SIDS, and may even

lower the risk. [67,69] One of the authors (VAS) presented a detailed analysis why case-control studies showed no association between SIDS and vaccines [71].

SIDS case definition in the cohort study [24] did not require an autopsy. Whether the 47 out of 109 (43%) non-autopsied post-DTP infant deaths were indeed SIDS will never be known. This study, however, was cited in 2005 as a “proof” of absence of association between SIDS and vaccines [72].

Epidemiological studies on SIDS and vaccines did not disprove that a small number of post-vaccination infant deaths in general population was classified as unexplained even though the infants either experienced adverse vaccine reactions before expiring, or had serious and potentially fatal pathological findings as described in VAERS reports. Such signs, symptoms, and findings cannot be regarded as terminal events of *coincidental* unexplained deaths.

This review and previous analysis of fatalities reported to VAERS [29] support suggestions of Kalokerinos [73] and Clemetson [74] to postpone vaccination of premature or sick infants and to supplement vitamin C around the time of vaccination in order to reduce the number of adverse events and to lower infant mortality.

## 6. Conclusion

In most of the reviewed cases, the history, the symptoms and or the pathologic findings did not support the classification of death as unexplained or SIDS. All cases of “SIDS” occurring shortly after vaccination should be reported to VAERS.

The risks of vaccinating premature, small or slightly ill infants should be weighed against the benefits of each vaccine, particularly in cases where the vaccine is not even needed.

The role of low serum ascorbate in SIDS should be further investigated particularly since the number of recommended pediatric vaccines is constantly increasing.

## References

- [1] Adelson L, Kinney ER. Sudden and unexpected death in infancy and childhood. *Pediatrics* 1956;17:663–99.
- [2] Fitzgibbons JP Jr., Nobrega FT, Ludwig J, Kurland LT, Harris LE. Sudden, unexpected, and unexplained death in infants. *Pediatrics* 1969;43: 980–8.
- [3] Wedgwood RJ, Benditt EP, ed. Sudden death in infants. Proceedings of the Conference on Causes of Sudden Death in Infants, Seattle, WA. Public Health Service Publication No 1412, National Institute of Child Health and Human Development, Bethesda, Maryland, 1964.
- [4] Bergman AB, Beckwith JB, Ray CG, eds. Sudden Infant Death Syndrome. Proceedings of the Second International Conference on Causes of Sudden Death in Infants, Seattle and London: University of Washington Press, 1970:18.
- [5] Bergman AB, Ray CG, Pomeroy MA, Wahl PW, Beckwith JB. Studies of the Sudden Infant Death Syndrome in King County, Washington. III. Epidemiology. *Pediatrics* 1972;49:860–70.
- [6] Willinger M, James LS, Catz C. Defining the sudden infant death syndrome (SIDS): deliberations of an expert panel convened by the National Institute of Child Health and Human Development. *Pediatr Pathol* 1991;11:677–84.
- [7] MacDorman MF, Rosenberg HM. Trends in infant mortality by cause of death and other characteristics, 1960-88. *National Center for Health Statistics. Vital Health Stat* 1993;20(20).
- [8] Kraus JF, Borhani NO. Postneonatal sudden unexpected death in California: a cohort study. *Am J Epidemiol* 1972;95:497–510.

- [9] Peterson DR. Evolution of the epidemiology of sudden infant death syndrome. *Epidemiol Rev* 1980;2:97–112.
- [10] CDC. NCHS. Compressed Mortality File 1979-1998. CDC Wonder Online Database, compiled from Compressed Mortality File CMF 1968-1988, Series 20, No. 2A, 2000 and CMF 1989-1998, Series 20, No. 2E, 2003. Compressed Mortality File 1999-2004. CDCWONDER On-Line Database, compiled from Compressed Mortality File 1999-2004 Series 20, No. 2J, 2007. Available online at <http://wonder.cdc.gov/mortSQL.html>.
- [11] Jones AM, Weston JT. The examination of the sudden infant death syndrome infant: Investigative and autopsy protocols. *J Forensic Sci* 1976; 21:833–41.
- [12] American Academy of Pediatrics, Division of State Government Affairs, 2005 State Legislation Report, Elk Grove Village, IL.
- [13] Death Scene Investigation Protocol. <http://www.californiasids.com/Universal/MainPage.cfm?p=45#1>
- [14] CDC. Guidelines for death scene investigation of sudden, unexplained infant deaths: recommendations of the interagency panel of sudden infant death syndrome. *MMWR Morb Mortal Wkly Rep* 1996;45(RR-10):1–22.
- [15] Tomashek KM, Shapiro-Mendoza CK, Wingo J. National Standard for Death-Scene Investigation of Sudden, Unexpected Infant Deaths in the United States. *Pediatrics* 2005;115:823.
- [16] Hauck F. Final report: National survey to evaluate use of the Sudden Unexplained Infant Death Investigation Report Form (SUIDIRF). Charlottesville, VA: University of Virginia Health System, 2001.
- [17] <http://www.cdc.gov/SIDS/SUID.htm>
- [18] Release of sudden, unexplained infant death investigation reporting form. *JAMA* 2006;295:2351.
- [19] Smith MH. National Childhood Vaccine Injury Compensation Act. *Pediatrics* 1988;82:264–9.
- [20] Chen RT, Rastogi SC, Mullen JR, Hayes SW, Cochi SL, Donlon JA *et al*. The Vaccine Adverse Event Reporting System (VAERS). *Vaccine* 1994; 12:542–50.
- [21] Ridgway D. Disputed Claims for pertussis vaccine injuries under the National Vaccine Injury Compensation Program. *J Investig Med* 1998;46: 168–74.
- [22] CD-ROM Series 20, 1985 Birth Cohort Linked Birth/Infant Death Data Sets 1995 Feb; 4. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [23] Hoffman HJ, Hunter JC, Damus K, Pakter J, Peterson DR, van Belle G *et al*. Diphtheria-tetanus-pertussis immunization and sudden infant death: results of the National Institute of Child Health and Human Development Cooperative Epidemiological Study of Sudden Infant Death Syndrome Risk Factors. *Pediatrics* 1987;79:598–611.
- [24] Griffin MR, Ray WA, Livengood JR, Schaffner W. Risk of sudden infant death syndrome after immunization with the diphtheria-tetanus-pertussis vaccine. *N Engl J Med* 1988;319:618–23.
- [25] Krous HF, Beckwith JB, Byard RW, Rognum TO, Bajanski T, Corey T *et al*. Sudden infant death syndrome and unclassified sudden infant deaths: a definitional and diagnostic approach. *Pediatrics* 2004;114:234–8.
- [26] CDC. Hepatitis B virus: a comprehensive strategy for eliminating transmission in the United States through universal childhood vaccination: Recommendations of the Immunization Practices Advisory Committee (ACIP). *MMWR Morb Mortal Wkly Rep* 1991;40(RR-13):1-19.
- [27] Kattwinkel J, Brooks J, Myerberg D. American Academy of Pediatrics AAP Task Force on infant positioning and SIDS: positioning and SIDS [published correction appears in *Pediatrics* 1992;90:264]. *Pediatrics* 1992; 89:1120–6.
- [28] Malloy MH, Freeman DH. Age at death, season, and day of death as indicators of the effect of the back to sleep program on sudden infant death syndrome in the United States, 1992-1999. *Arch Pediatr Adolesc Med* 2004;158:359–65.
- [29] Silvers LE, Ellenberg SS, Wise RP, Varricchio FE, Mootrey GT, Salive ME. The epidemiology of fatalities reported to the Vaccine Adverse Event Reporting System 1990-1997. *Pharmacoepidemiology and Drug Safety* 2001;10:279–85.
- [30] <http://www.medalerts.org> (last accessed on February 22, 2007)
- [31] CD-ROM Series 20, 1992 Multiple Cause-of-Death File 1997 Jun; 10. United States Department of Health and Human Services (US DHHS), Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Hyattsville, Maryland.
- [32] CD-ROM Series 20, 1993 Multiple Cause-of-Death File 1997 Jul; 11. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [33] CD-ROM Series 20, 1994 Multiple Cause-of-Death File 1999 Nov; 13. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [34] CD-ROM Series 20, 1995 Perinatal Mortality Data File 1998 Apr; 12. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [35] CD-ROM Series 20, 1996 Perinatal Mortality Data File 1998 Nov; 14. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [36] CD-ROM Series 20, 1997 Perinatal Mortality Data File 1999 Sep; 15. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [37] CD-ROM Series 20, 1998 Perinatal Mortality Data File 2000 Dec; 18. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [38] CD-ROM Series 20, 1999 Perinatal Mortality Data File 2002 Apr; 20. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [39] CD-ROM Series 20, 2000 Perinatal Mortality Data File 2003 Feb; 21. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [40] CD-ROM Series 20, 2001 Perinatal Mortality Data File 2004 Feb; 22. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [41] CD-ROM Series 20, 2002 Perinatal Mortality Data File 2005 Jan; 23. US DHHS, CDC, NCHS, Hyattsville, Maryland.
- [42] Niu MT, Salive ME, Ellenberg SS. Neonatal deaths after hepatitis B vaccine: the vaccine adverse event reporting system, 1991-1998. *Arch Pediatr Adolesc Med* 1999;153:1279–82.
- [43] MacDorman MF, Atkinson JO. Infant mortality statistics from the 1997 period linked birth/infant death data set. National vital statistics reports 1999;47(23). Hyattsville, Maryland: National Center for Health Statistics.
- [44] <http://www.childdeathreview.org/spotlightNH.htm> New Hampshire Child Fatality Review Committee Fourth Annual Report 2001 Nov: 6.
- [45] Centers for Disease Control and Prevention. *Epidemiology and Prevention of Vaccine-Preventable Diseases*. Atkinson W, Hamborsky J, McIntyre L, Wolfe S, eds. 9th ed. Washington DC: Public Health Foundation, 2006.
- [46] Valdés-Dapena MA, McFeeley PA, Hoffman HJ, Damus KH, Franciosi RR, Allison DJ *et al*. Histopathology atlas for the Sudden Infant Death Syndrome. Washington, D.C.: Armed Forces Institute of Pathology in cooperation with the American Registry of Pathology and the National Institute of Child Health and Human Development, 1993.
- [47] Goldwater PN. Sudden infant death syndrome: a critical review of approaches to research. *Arch Dis Child* 2003;88:1095–100.
- [48] Kalokerinos A, Dettman G. Sudden death in infancy syndrome in Western Australia *Med J Aust* 1976;2:31–2.
- [49] Bousquet J, Chanez P, Lacoste JY, Barnéon G, Ghavanian N, Enander I *et al*. Eosinophilic inflammation in asthma. *N Engl J Med* 1990;323:1033–9.
- [50] Howat WJ, Moore IE, Judd M, Roche WR. Pulmonary immunopathology of sudden infant death syndrome. *Lancet* 1994;343:1390–2.
- [51] Osborn GR. Findings in 262 fatal accidents. *Lancet* 1943;2:277.
- [52] Follis RH. Sudden death in infants with scurvy. *J Pediatr* 1942;20:347–51.
- [53] Kalokerinos A. Every Second Child. New Canaan, Conn.: Keats Publishing, 1981.
- [54] Cheraskin, E. Letter to the Editor: Vitamin C, smoking and SIDS. *J R Soc Health* 1995;115:332.
- [55] Kalokerinos A. The sudden infant death syndrome. Part 2. Definition. Further clinical observations. *Australas Nurses J* 1978;7:6–8.
- [56] NCHS. Health, United States, 1993. Hyattsville, MD: Public Health Service, 1994.
- [57] Kramarow E, Lentzner H, Rooks R, Weeks J, Saydah S. Health and Aging Chartbook. Health, United States, 1999. Hyattsville, Maryland: National Center for Health Statistics, 1999.
- [58] Botham SJ, Isaacs D. Incidence of apnoea and bradycardia in preterm infants following triple antigen immunization. *J Paediatr Child Health* 1994;30:533–5.
- [59] Botham SJ, Isaacs D, Henderson-Smart DJ. Incidence of apnoea and bradycardia in preterm infants following DTPw and Hib immunization: a prospective study. *J Paediatr Child Health* 1997; 33:418–21.
- [60] Slack MH, Schapira D. Severe apnoeas following immunisation in premature infants. *Arch Dis Child Fetal Neonatal Ed* 1999;81:F67–8.
- [61] Sen S, Cloete Y, Hassan K, Buss P. Adverse events following vaccination in premature infants. *Acta Paediatr* 2001;90:916–20.
- [62] Sanchez PJ, Laptook AR, Fisher L, Sumner J, Risser RC, Perlman JM. Apnea after immunization of preterm infants. *J Pediatr* 1997;130:746–51.
- [63] [http://www.cdc.gov/od/science/iso/concerns/archive/sids\\_faq.htm](http://www.cdc.gov/od/science/iso/concerns/archive/sids_faq.htm)
- [64] Malloy MH, MacDorman M. Changes in the classification of sudden unexpected infant deaths: United States, 1992-2001. *Pediatrics*. 2005;115: 1247-1253.
- [65] Shapiro-Mendoza CK, Tomashek KM, Anderson RN, Wingo J. Recent national trends in sudden, unexpected infant deaths: more evidence supporting a change in classification or reporting. *Am J Epidemiol*. 2006;163: 762–9.

- [66] MacDorman MF, Martin JA, Mathews TJ, Hoyert DL, Ventura SJ. Explaining the 2001–02 infant mortality increase: Data from the linked birth/infant death data set. National vital statistics reports 2005;53(12). Hyattsville, Maryland: National Center for Health Statistics.
- [67] Mitchell EA, Stewart AW, Clements M, Ford RPK. Immunisation and the sudden infant deaths syndrome. Arch Dis Child 1995; 73: 498–501.
- [68] Jonville-Bera AP, Autret-Leca E, Barbeillon F, Paris-Llado J; French Reference Centers for SIDS. Sudden unexpected death in infants under 3 months of age and vaccination status—a case-control study. Br J Clin Pharmacol 2001;51:271–6.
- [69] Fleming PJ, Blair PS, Platt MW, Tripp J, Smith IJ, Golding J. The UK accelerated immunisation programme and sudden unexpected death in infancy: case-control study. BMJ 2001;322:822–25. Available online at <http://www.bmj.com/cgi/content/full/322/7290/822>
- [70] Vennemann MMT, Butterfass-Bahloul T, Jorch G, Brinkmann B, Findeisen M, Sauerland C *et al.* Sudden infant death syndrome: No increased risk after immunization. Vaccine 2007;25:336–40.
- [71] Soldatenkova VA. Why case-control studies showed no association between Sudden Infant Death Syndrome and vaccinations. Medical Veritas 2007 Apr.;4(1):1411–3.
- [72] American Academy of Pediatrics, Task Force on Sudden Infant Death Syndrome. The changing concept of sudden infant death syndrome: diagnostic coding shifts, controversies regarding the sleeping environment, and new variables to consider in reducing risk. Pediatrics. 2005;116: 1245–55.
- [73] Kalokerinos A. Medical Pioneer of the 20<sup>th</sup> Century. An Autobiography. Braeside, Melbourne, Victoria: Biological Therapies Publishing Pty Ltd, 2000.
- [74] Clemetson CAB. Vaccinations, Inoculations and Ascorbic Acid. J Orthomol Med 1999;14:137–42.

**Table 1. Unexplained neonatal deaths following hepatitis B vaccination given alone with corresponding vaccine lot numbers reported to VAERS, Jan. 1992 – Dec. 2002**

| Case no. | VAERS ID           | Sex | State | Lot number | Number of reports <sup>b</sup> | Vaccination date | Days since vaccination | Age at death (days) |
|----------|--------------------|-----|-------|------------|--------------------------------|------------------|------------------------|---------------------|
| 1        | 46348 <sup>a</sup> | M   | VA    | 1032A4     | 1                              | 10/08/92         | 2                      | 15                  |
| 2        | 51067              | M   | PA    | 1116A2     | 9 (3)                          | 03/15/93         | 1                      | 28                  |
| 3        | 51688              | F   | PA    | 116A2      | 1                              | 03/29/93         | 2                      | 15                  |
| 4        | 51976              | F   | SC    | 01554V     | 1                              | 03/17/93         | 14                     | 30 <sup>c</sup>     |
| 5        | 51982              | M   | PA    | 1112A2     | 10 (2)                         | 04/02/93         | 1                      | 30 <sup>c</sup>     |
| 6        | 52816              | F   | MD    | 1119A2     | 6 (1)                          | 05/14/93         | 2                      | 6                   |
| 7        | 59322              | F   | SC    | 0942W      | 32 (2)                         | 11/30/93         | 10                     | 24                  |
| 8        | 70603              | F   | NH    | 1547W      | 32 (3)                         | 06/24/94         | 19                     | 22                  |
| 9        | 87649              | M   | OR    | 1189B      | 15 (2)                         | 06/28/96         | 2                      | 14                  |
| 10       | 91425              | M   | NH    |            |                                | 09/05/96         | 21                     | 24                  |
| 11       | 97016              | M   | NH    | 0165D      | 14 (3)                         | 09/02/96         | 54                     | 55                  |
| 12       | 108796             | M   | NH    |            |                                | 01/02/97         | 18                     | 19                  |
| 13       | 96889              | F   | NH    | 1289A      | 18 (3)                         | 09/24/96         | 9                      | 22                  |
| 14       | 107414             | F   | OH    | 1611B      | 22 (1)                         | 04/3/96          | 38                     | 39                  |
| 15       | 108795             | F   | NH    | 1163A      | 28 (1)                         | 04/13/96         | 15                     | 16                  |
| 16       | 112250             | F   | NY    | 2587A2     | 15 (1)                         | 06/15/98         | <1                     | 21                  |
| 17       | 113478             | M   | NH    | 0424E      | 1                              | 11/24/97         | 32                     | 34                  |
| 18       | 117475             | M   | NH    | 1559E      | 9 (1)                          | 10/24/98         | 25                     | 27                  |
| 19       | 118117             | M   | WV    | 2612A2     | 28 (1)                         | 05/29/98         | 1                      | 19                  |
| 20       | 122713             | M   | NH    |            |                                | 03/3/99          | 18                     | 20                  |
| 21       | 123168             | M   | NH    | 2930A2     | 20 (3)                         | 03/24/99         | 31                     | 32                  |
| 22       | 126759             | M   | NH    | 0987H      | 13 (3)                         | 11/07/98         | 11                     | 13                  |
| 23       | 133751             | M   | NH    | 2948A2     | 24 (1)                         | 05/27/99         | 16                     | 17                  |
| 24       | 171973             | F   | NH    | 1947J      | 3 (1)                          | 05/7/00          | 20                     | 21                  |
| 25       | 171370             | F   | NH    | ENG518A2   | 4 (2)                          | 02/26/01         | 6                      | 26                  |
| 26       | 200120             | M   | NH    | 5286A2     | 26 (1)                         | 12/19/02         | 5                      | 7                   |

<sup>a</sup> Reported age at death was 15 days [42], but no such corresponding death record from Virginia was found in the 1992 mortality file [31].

<sup>b</sup> Number of death reports from each lot is listed in the parentheses.

<sup>c</sup> Age at death is 1 month in 1993 main mortality file [32].

**Table 2. Lot Number not reported**

| Case No. | VAERS ID | Sex | State | Vaccine date | Days since vaccination | Age at death (days) |
|----------|----------|-----|-------|--------------|------------------------|---------------------|
| 27       | 100366   | M   | MS    | 070/6/97     | 4                      | 20                  |
| 28       | 150249   | F   | NJ    | 02/28/00     | 4                      | 11                  |
| 29       | 182530   | M   | IN    | 10/29/01     | 3                      | 5                   |

**Table 3. Cases of unexplained infant deaths, New Hampshire, U.S., 1979-1990 (Underlying cause of death codes 798.0 and 799.9 respectively, ICD-9)**

| Year | Live births | Neonatal cases |  | Postneonatal cases |  | Total |
|------|-------------|----------------|--|--------------------|--|-------|
|      |             | SIDS and 799.9 |  | SIDS and 799.9     |  |       |
| 1979 | 12,845      | 1              |  | 16                 |  | 17    |
| 1980 | 13,745      | 1              |  | 12                 |  | 13    |
| 1981 | 13,517      | 2              |  | 22                 |  | 24    |
| 1982 | 14,087      | 0              |  | 9                  |  | 9     |
| 1983 | 13,799      | 3 and 1        |  | 17 and 1           |  | 22    |
| 1984 | 14,250      | 2              |  | 15 and 1           |  | 18    |
| 1985 | 15,453      | 1              |  | 19                 |  | 20    |
| 1986 | 15,895      | 4              |  | 15                 |  | 19    |
| 1987 | 17,032      | 5              |  | 15 and 1           |  | 21    |
| 1988 | 17,364      | 1 and 1        |  | 25 and 2           |  | 29    |
| 1989 | 17,809      | 1              |  | 26 and 1           |  | 28    |
| 1990 | 17,569      | 0              |  | 18 and 4           |  | 22    |

**Table 4. Cases of unexplained infant deaths (UID), New Hampshire, U.S., 1991-2002 (ICD-9 codes are 798.0 and 799.9 for 1991-1998; R95 and R99 respectively since 1999, ICD-10)**

| Year | Live births | Neonatal Cases     |                   | Postneonatal Cases |                   |           |                   | Total UID reported to VAERS |       |
|------|-------------|--------------------|-------------------|--------------------|-------------------|-----------|-------------------|-----------------------------|-------|
|      |             | SIDS and 799.9/R99 |                   | SIDS               |                   | 799.9/R99 |                   | VAERS                       | Total |
|      |             | All                | Reported to VAERS | All                | Reported to VAERS | All       | Reported to VAERS |                             |       |
| 1991 | 16,341      | 2                  | 0                 | 14                 | 2                 | 4         | 1                 | 3                           | 20    |
| 1992 | 15,990      | 1                  | 0                 | 13                 | 2                 | 3         | 1                 | 3                           | 17    |
| 1993 | 15,436      | 0                  | 0                 | 15                 | 4                 | 1         | 0                 | 4                           | 16    |
| 1994 | 15,106      | 1                  | 1                 | 14                 | 4                 | 3         | 1                 | 6                           | 18    |
| 1995 | 14,665      | 3                  | 0                 | 12                 | 3                 | 1         | 1                 | 4                           | 16    |
| 1996 | 14,520      | 4                  | 3                 | 5                  | 2                 | 0         | 0                 | 5                           | 9     |
| 1997 | 14,313      | 1                  | 1                 | 4                  | 4                 | 3         | 2                 | 7                           | 8     |
| 1998 | 14,429      | 4                  | 2                 | 4                  | 2                 | 0         | 0                 | 4                           | 8     |
| 1999 | 14,041      | 2                  | 2                 | 3                  | 2                 | 0         | 0                 | 4                           | 5     |
| 2000 | 14,609      | 3                  | 1                 | 4                  | 3                 | 5         | 2                 | 6                           | 12    |
| 2001 | 14,656      | 1                  | 1                 | 7                  | 0                 | 0         | 0                 | 1                           | 8     |
| 2002 | 14,442      | 3                  | 1                 | 6                  | 2                 | 0         | 0                 | 3                           | 8     |

**Table 5. U.S. Resident Births and Infant Deaths, 1989-2004**

| Year | No. of births | Death Counts |             |            |             |
|------|---------------|--------------|-------------|------------|-------------|
|      |               | SIDS         |             | All Causes |             |
|      |               | 0-27 Days    | 28-364 Days | 0-27 Days  | 28-364 Days |
| 1989 | 4,040,958     | 398          | 5,236       | 25,168     | 14,487      |
| 1990 | 4,158,212     | 367          | 5,050       | 24,309     | 14,042      |
| 1991 | 4,110,907     | 286          | 5,063       | 22,978     | 13,788      |
| 1992 | 4,065,014     | 298          | 4,593       | 21,849     | 12,779      |
| 1993 | 4,000,240     | 287          | 4,382       | 21,174     | 12,292      |
| 1994 | 3,952,767     | 258          | 3,815       | 20,250     | 11,460      |
| 1995 | 3,899,589     | 219          | 3,178       | 19,155     | 10,428      |
| 1996 | 3,891,494     | 213          | 2,837       | 18,572     | 9,915       |
| 1997 | 3,880,894     | 182          | 2,809       | 18,524     | 9,521       |
| 1998 | 3,941,553     | 188          | 2,634       | 18,918     | 9,453       |
| 1999 | 3,959,417     | 208          | 2,440       | 18,728     | 9,209       |
| 2000 | 4,058,814     | 204          | 2,319       | 18,776     | 9,259       |
| 2001 | 4,025,933     | 187          | 2,047       | 18,265     | 9,303       |
| 2002 | 4,021,726     | 185          | 2,110       | 18,747     | 9,287       |
| 2003 | 4,089,950     | 190          | 1,972       | 18,893     | 9,132       |
| 2004 | 4,112,052     | 210          | 2,036       | 18,593     | 9,343       |

**Table 6. Neonatal and Postneonatal Mortality Rates for SIDS and All Causes per 1,000 Live Births, 1989-2004**

| Year | SIDS Mortality Rates |                        | All Causes Mortality Rates |                        |
|------|----------------------|------------------------|----------------------------|------------------------|
|      | 0-27 Days (95% C.I.) | 28-364 Days (95% C.I.) | 0-27 Days (95% C.I.)       | 28-364 Days (95% C.I.) |
| 1989 | 0.098 (0.089, 0.108) | 1.296 (1.260, 1.331)   | 6.228 (6.151, 6.305)       | 3.585 (3.527, 3.644)   |
| 1990 | 0.088 (0.079, 0.097) | 1.214 (1.181, 1.248)   | 5.846 (5.772, 5.920)       | 3.377 (3.321, 3.433)   |
| 1991 | 0.070 (0.062, 0.078) | 1.231 (1.198, 1.266)   | 5.590 (5.517, 5.662)       | 3.354 (3.298, 3.410)   |
| 1992 | 0.073 (0.065, 0.082) | 1.130 (1.097, 1.163)   | 5.375 (5.303, 5.446)       | 3.144 (3.089, 3.198)   |
| 1993 | 0.072 (0.063, 0.080) | 1.095 (1.063, 1.128)   | 5.293 (5.221, 5.365)       | 3.073 (3.018, 3.127)   |
| 1994 | 0.065 (0.057, 0.073) | 0.965 (0.935, 0.996)   | 5.123 (5.052, 5.194)       | 2.900 (2.846, 2.952)   |
| 1995 | 0.056 (0.049, 0.064) | 0.815 (0.787, 0.843)   | 4.912 (4.842, 4.982)       | 2.674 (2.623, 2.726)   |
| 1996 | 0.055 (0.047, 0.062) | 0.729 (0.702, 0.756)   | 4.772 (4.704, 4.841)       | 2.548 (2.498, 2.598)   |
| 1997 | 0.047 (0.040, 0.054) | 0.724 (0.697, 0.751)   | 4.773 (4.704, 4.842)       | 2.453 (2.404, 2.503)   |
| 1998 | 0.048 (0.041, 0.055) | 0.668 (0.643, 0.694)   | 4.800 (4.731, 4.868)       | 2.398 (2.350, 2.447)   |
| 1999 | 0.052 (0.045, 0.060) | 0.616 (0.592, 0.641)   | 4.730 (4.662, 4.798)       | 2.326 (2.278, 2.373)   |
| 2000 | 0.050 (0.043, 0.057) | 0.571 (0.548, 0.595)   | 4.626 (4.560, 4.692)       | 2.281 (2.235, 2.328)   |
| 2001 | 0.046 (0.040, 0.053) | 0.508 (0.486, 0.530)   | 4.537 (4.471, 4.603)       | 2.311 (2.264, 2.358)   |
| 2002 | 0.046 (0.039, 0.053) | 0.525 (0.502, 0.547)   | 4.661 (4.595, 4.728)       | 2.309 (2.262, 2.356)   |
| 2003 | 0.046 (0.040, 0.053) | 0.482 (0.461, 0.503)   | 4.619 (4.553, 4.685)       | 2.233 (2.187, 2.279)   |
| 2004 | 0.051 (0.044, 0.058) | 0.495 (0.474, 0.517)   | 4.522 (4.456, 4.587)       | 2.272 (2.226, 2.318)   |